

HORSE FIOC H 8 puledri

12/36

COMPLEMENTARY FEED FOR HORSES

Additives per Kg

Vitamins, pro-vitamins and chemically well-defined substances having similar effect

3a672a Vitamin A (Retinyl Acetate)	12,000,00 UI
3a671 Vitamin D3 (cholecalciferol)	1,200,00 UI
3a880 Biotin	0,24 mg
3a890 Choline chloride	390,00 mg
3a316 Folic acid	8,00 mg
3a315 Niacinamide	13,50 mg
3a841 Calcium-D-pantothenate	21,25 mg
3a821 Vitamin B1 (Thiamine mononitrate)	10,50 mg
3a837 Vitamin B12 (Cyanocobalamin)	0,10 mg
3a825i Vitamin B2 (Riboflavin)	16,95 mg
3a831 Vitamin B6 (pyridoxine hydrochloride)	5,00 mg
3a300 Vitamin C (Ascorbic acid)	200,00 mg
3a700 Vitamin E (all-rac-alpha-tocopheryl acetate)	84,00 mg
3a711 Vitamin K3 (Menadione nicotinamide bisulphite)	8,00 mg

Compounds of trace elements

3b407 Copper (Copper -II- chelate of protein hydrolysates)	5,00 mg
3b405 Copper (Copper -II- sulphate pentahydrate)	45,00 mg
3b201 Iodine (Potassium iodide)	0,45 mg
3b101 Iron (Iron -II- carbonate siderite)	102,60 mg
3b107 Iron (Iron -II- chelate of protein hydrolysates)	11,40 mg
3b502 Manganese (manganese II oxide)	121,50 mg
3b505 Manganese (manganese chelate of protein hydrolysates)	13,50 mg
3b801 Selenium (Sodium selenite)	0,24 mg
3b603 Zinc (Zinc oxide)	135,00 mg
3b612 Zinc (Zinc chelate of protein hydrolysates)	15,00 mg

Amino acids, their salts and analogues

3c322 L-lysine monohydrochloride, technically pure	1.560,00 mg
--	-------------

Antioxidants

E 321 Butylated hydroxytoluene (BHT)	60,00 mg
--------------------------------------	----------

COMPOSITION

Whole barley flakes, Wheat bran, Corn flake, Oats, Beet pulp, Whole soy flake(*), Flour extraction of dehulled sunflower seeds, Dried Alfalfa, Rice husk, Molasses and soy oil, Flaked Fava bean, Whole lin seed, Calcium carbonate, Sodium chloride, Soy oil, Sweet dried serum, Magnesium oxyde, Zeolyte, Bicalic phosphate.

ANALITICAL COMPONENTS

Crude protein	13,40 %
Crude oils and fats	5,80 %
Crude fiber	8,00 %
Crude ash	8,00 %
Sodium	0,68 %

FEEDING USE

HORSE FIOC H8 must be used for foals from the 12th up to the 36th month of birth. Dosage: 1% of its body weight. If the animal is subjected to prolonged physical effort the ration may be increased during this period from 1.2 to 1.5% depending to weight and behavior of the animal. This product is optimal for stallions also. Leave hay and clean water at free disposal to the horse.



To be consumed within:

LOT Nr.: 202608311050 02/2027

The chemical analysis and vitamin content are guaranteed only if the product is stored in a cool and dry place and if sealed in its original packaging. Produced in the plant with ministerial authorization n. MG/143-R/A 1310 del 06/10/1982 alfa IT 000187 TV